


Recall: Yesterday we learned that if a triangle is a right triangle, and you know two of its side lengths, then you can find the third using the Pythagorean Theorem. However, could we use the Pythagorean Theorem to determine if a triangle is a right triangle or not? This is called the converse of the Pythagorean Theorem.

The Converse of the Pythagorean Theorem

If the square of the length of the longest side of a triangle is equal to the sum of the square of the lengths of the other two sides, then the triangle is a right triangle.

If  $a^2 + b^2 = c^2$ , then  $\triangle ABC$  is a right triangle.



Do the given side lengths form a right triangle? Explain why or why not. (Remember, the hypotenuse will always be the longest length)

$\approx 2.2$      $5.6$   
 1.  $\sqrt{5}, 6, \sqrt{31}$

$$(\sqrt{5})^2 + (\sqrt{31})^2 \stackrel{?}{=} 6^2$$

$$5 + 31 \stackrel{?}{=} 36$$

$$36 = 36 \quad \text{YES}$$

2. 33, 20, 28

$$20^2 + 28^2 \stackrel{?}{=} 33^2$$

$$400 + 784 \stackrel{?}{=} 1089$$

$$1184 \neq 1089$$

NO

3. 25, 7, 24

$$7^2 + 24^2 \stackrel{?}{=} 25^2$$

$$49 + 576 \stackrel{?}{=} 625$$

$$625 = 625 \quad \text{YES}$$

$\approx 4.2$   
 4. 13,  $3\sqrt{2}$ , 9

$$(3\sqrt{2})^2 + 9^2 \stackrel{?}{=} 13^2$$

$$(9)(2) + 81 \stackrel{?}{=} 169$$


$$18 + 81 \stackrel{?}{=} 169$$

$$99 \neq 169$$

NO

If the sides of a triangle do not form right triangles, then there are two other options. Either they could be *acute triangles* or they could be *obtuse triangles*. It turns out that when numbers are substituted into the Pythagorean Theorem, there is a way to tell if the triangle will be acute or obtuse.

Classifying Triangles



In  $\triangle ABC$  with the longest side  $c$ :

If  $a^2 + b^2 > c^2$ , then  $\triangle ABC$  is an **ACUTE** triangle.

>

**ACUTE**

If  $a^2 + b^2 = c^2$ , then  $\triangle ABC$  is a **RIGHT** triangle.

=


**RIGHT**

If  $a^2 + b^2 < c^2$ , then  $\triangle ABC$  is an **OBTUSE** triangle.

<

**OBTUSE**

Classify each triangle as acute, right, or obtuse.




$$4^2 + 5^2 = (\sqrt{35})^2$$

$$16 + 25 \stackrel{?}{=} 35$$

$$41 > 35$$

Acute




$$12^2 + 16^2 \stackrel{?}{=} 20^2$$

$$144 + 256 \stackrel{?}{=} 400$$

$$400 = 400$$

Right




$$15^2 + 17^2 \stackrel{?}{=} 23^2$$

$$225 + 289 \stackrel{?}{=} 529$$

$$514 < 529$$

Obtuse




$$8^2 + 8^2 \stackrel{?}{=} 12^2$$

$$64 + 64 \stackrel{?}{=} 144$$

$$128 < 144$$

obtuse

9. A steel pole 150 cm in length has been placed in the ground ready for cement to seal it in place. To check to see if it is perpendicular to the ground, the contractor has measured a distance of 180 cm from the top of the pole to 80 cm from the base of the pole on the ground. Is the pole perpendicular to the ground?




$$80^2 + 150^2 \stackrel{?}{=} 180^2$$

$$6400 + 22500 \stackrel{?}{=} 32400$$

NO

Review

10. If the two legs of a right triangle are 4 and 6, what is the length of the hypotenuse? Keep your answer in simplest radical form.



$$4^2 + 6^2 = c^2$$

$$16 + 36 = c^2$$

$$52 = c^2$$

$$\sqrt{52}$$

$$2 \sqrt{13}$$